

ChronoBiology Letter #18 Article References Salivary Cortisol - A Biomarker for Depression

by: John J. White, MD CM

1. Herbert J. Cortisol and depression: three questions for psychiatry. *Psychol Med.* 2013;43(3):449-69.
2. Holsboer F. The corticosteroid receptor hypothesis of depression *Neuropsychopharmacology.* 2000;23(5):477-501.
3. Stetler C, Miller GE. Depression and hypothalamic-pituitary-adrenal activation: a quantitative summary of four decades of research. *Psychosom Med.* 2011;73(2):114-26.
4. Guerry JD, Hastings PD. In search of HPA axis dysregulation in child and adolescent depression. *Clin Child Fam Psychol Rev.* 2011;14(2):135-60.
5. Goodyer IM, Bacon A, Ban M, Croudace T, Herbert J. Serotonin transporter genotype, morning cortisol and subsequent depression in adolescents. *Br J Psychiatry.* 2009;195(1):39-45.
6. Goodyer IM, Herbert J, Tamplin A, Altham PM. First-episode major depression in adolescents: affective, cognitive and endocrine characteristics of risk status and predictors of onset. *Br J Psychiatry.* 2000;176:142-9.
7. Owens M, Herbert J, Jones PB, et al. Elevated morning cortisol is a stratified population-level biomarker for major depression in boys only with high depressive symptoms. *Proc Natl Acad Sci U S A.* 2014;111(9):3638-43.
8. DeGroot L, Jameson J. *Endocrinology.* Vol 1. 5th ed. Philadelphia, PA: Elsevier Saunders; 2006:438.



ChronoBiology Letter #18 Article References

Amoebiasis

by: Kamal Henein, MD

1. Ali IK, Clark CG, Petri WA, Jr. Molecular epidemiology of amebiasis. *Infect Genet Evol.* 2008;8(5):698-707.
2. Van Hal SJ, Stark DJ, Fotedar R, Marriot D, Ellis JT, Harkness JL. Amoebiasis: current status in Australia. *Med J Aust.* 2007;186(8):412-6.
3. Ximénez C, Morán P, Rojas L, Valadez A, Gomez A. Reassessment of the epidemiology of amebiasis: state of the art. *Infect Genet Evol.* 2009;9(6):1023-32.
4. Chacín-Bonilla L. Amebiasis: implications of the recognition of *Entamoeba dispar* and the identification of *Entamoeba moshkovskii* in humans. *Invest Clin.* 2010;51(2):239-56.
5. Stanley SL, Jr. Amoebiasis. *Lancet.* 2003;361(9362):1025-34.
6. Fotedar R, Stark D, Beebe N, Marriot D, Ellis J, Harkness J. Laboratory diagnostic techniques for *Entamoeba* species. *Clin Microbiol Rev.* 2007;20(3):511-32.
7. Walderich B, Weber A, Knobloch J. Differentiation of *Entamoeba histolytica* and *Entamoeba dispar* from German travelers and residents of endemic areas. *Am J Trop Med Hyg.* 1997;57(1):70-4.
8. Herbinger KH, Fleischmann E, Weber C, et al. Epidemiological, clinical, and diagnostic data on intestinal infections with *Entamoeba histolytica* and *Entamoeba dispar* among returning travelers. *Infection.* 2011;39(6):527-35.
9. Freedman DO, Weld LH, Kozarsky PE, et al. Spectrum of disease and relation to place of exposure among ill returned travelers. *N Engl J Med.* 2006;354(2):119-30.
10. Salit IE, Khairnar K, Gough K, Pillai DR. A possible cluster of sexually transmitted *Entamoeba histolytica*: genetic analysis of a highly virulent strain. *Clin Infect Dis.* 2009;49(3):346-53.



ChronoBiology Letter #18 Article References

Amoebiasis, cont.

11. Gunther J, Shafir S, Bristow B, Sorvillo F. Short report: Amebiasis-related mortality among United States residents, 1990-2007. *Am J Trop Med Hyg.* 2011;85(6):1038-40.
12. Hung CC, Ji DD, Sun HY, et al. Increased risk for *Entamoeba histolytica* infection and invasive amebiasis in HIV seropositive men who have sex with men in Taiwan. *PLoS Negl Trop Dis.* 2008;2(2):e175.
13. Hung CC, Chang SY, Ji DD. *Entamoeba histolytica* infection in men who have sex with men. *Lancet Infect Dis.* 2012;12(9):729-36.
14. Okamoto M, Kawabe T, Ohata K, et al. Amebic colitis in asymptomatic subjects with positive fecal occult blood test results: clinical features different from symptomatic cases. *Am J Trop Med Hyg.* 2005;73(5):934-5.
15. Verkerke HP, Petri WA Jr, Marie CS. The dynamic interdependence of amebiasis, innate immunity, and undernutrition. *Semin Immunopathol.* 2012;34(6):771-85.
16. Ravdin JI, Stanley P, Murphy CF, Petri WA, Jr. Characterization of cell surface carbohydrate receptors for *Entamoeba histolytica* adherence lectin. *Infect Immun.* 1989;57(7):2179-86.
17. Rayan HZ. Microscopic overdiagnosis of intestinal amoebiasis. *J Egypt Soc Parasitol.* 2005; 35(3):941-51.
18. Abd-Alla MD, Jackson TF, Gathiram V, el-Hawey AM, Ravdin JI. Differentiation of pathogenic *Entamoeba histolytica* infections from nonpathogenic infections by detection of galactose-inhibitable adherence protein antigen in sera and feces. *J Clin Microbiol.* 1993;31(11):2845-50.
19. Kimura M, Nakamura T, Nawa Y. Experience with intravenous metronidazole to treat moderate-to-severe amebiasis in Japan. *Am J Trop Med Hyg.* 2007;77(2):381-5.
20. Misra NP, Gupta RC. A comparison of a short course of single daily dosage therapy of tinidazole with metronidazole in intestinal amoebiasis. *J Int Med Res.* 1977;5(6):434-7.



ChronoBiology Letter #18 Article References Amoebiasis, cont.

21. Singh G, Kumar S. Short course of single daily dosage treatment with tinidazole and metronidazole in intestinal amoebiasis: a comparative study. *Curr Med Res Opin.* 1977;5(2):157-60.
22. Rossignol JF, Kabil SM, El-Gohary Y, Younis AM. Nitazoxanide in the treatment of amoebiasis. *Trans R Soc Trop Med Hyg.* 2007;101(10):1025-31.
23. Li E, Stanley SL, Jr. Protozoa. Amebiasis. *Gastroenterol Clin North Am.* 1996;25(3):471-92.
24. Mansour-Ghanaei F, Dehbashi N, Yazdanparast K, Shafaghi A. Efficacy of *Saccharomyces boulardii* with antibiotics in acute amoebiasis. *World J Gastroenterol.* 2003;9(8):1832-3.
25. Haque R, Duggal P, Ali IM, et al. Innate and acquired resistance to amebiasis in Bangladeshi children. *J Infect Dis.* 2002;186(4):547-52.

Figure 1: Microscopic view of *E. histolytica* trophozoite. Photo courtesy of CDC website.
http://www.cdc.gov/dpdx/images/intestinalAmebae/Ecoli_troph_tric.jpg

Figure 2: Copyright 2014 Diagnos-Techs, Inc.



ChronoBiology Letter #18 Article References Dr. Dialogue

Testosterone:

1. Finkle WD, Greenland S, Ridgeway GK, et al. Increased risk of non-fatal myocardial infarction following testosterone therapy prescription in men. *PLoS One*. 2014;9(1):e85805. doi: 10.1371/journal.pone.0085805.
2. Yeap BB, Alfonso H, Chubb SA, et al. In older men an optimal plasma testosterone is associated with reduced all-cause mortality and higher dihydrotestosterone with reduced ischemic heart disease mortality, while estradiol levels do not predict mortality. *J Clin Endocrinol Metab*. 2014;99(1):E9-18. doi: 10.1210/jc.2013-3272.

Vitamin D:

3. Vogiatzi MG, Jacobson-Dickman E, DeBoer MD, et al. Vitamin D supplementation and risk of toxicity in pediatrics: a review of current literature. *J Clin Endocrinol Metab*. 2014;99(4):1132-41. doi: 10.1210/jc.2013-3655. *Anticancer Res*. 2014 Mar;34(3):1163-6
4. Mohr SB, Gorham ED, Kim J, Hofflich H, Garland CF. Meta-analysis of vitamin D sufficiency for improving survival of patients with breast cancer. *Anticancer Res*. 2014 Mar;34(3):1163-6.

Effect of Chronic Stress on Children:

5. Shalev I, Moffitt TE, Sugden K, et al. Exposure to violence during childhood is associated with telomere erosion from 5 to 10 years of age: a longitudinal study. *Mol Psychiatry*. 2013 May;18(5):576-81. doi: 10.1038/mp.2012.32.
6. Mitchell C, Hobcraft J, McLanahan SS, et al. Social disadvantage, genetic sensitivity, and children's telomere length. *Proc Natl Acad Sci U S A*. 2014;111(16):5944-9. doi: 10.1073/pnas.1404293111.

